

Instructions: You have a total of 50 minutes to complete this test.

Answer each question completely showing complete details.

For complete credit you must include correct SI units with numerical answers.

Time Start _____ Time finish _____ pledged _____

$$\text{Constants: } g = 9.8 \frac{\text{m}}{\text{s}^2}$$

[1] A particle is observed to move with an acceleration given by:

$$\vec{a} = 0\hat{x} + ct^2\hat{y}$$

where the constant b has SI units of $[c] = \left[\frac{\text{m}}{\text{s}^4}\right]$.

(a) Find the velocity **vector** at a later time assuming at $t=0$ the vector velocity is zero.

(b) Find the position **vector** at a later time assuming at $t=0$, the velocity and position are both zero.

Suppose another particle is seen to have a position vector given by:

$$\vec{R}(t) = (-ct^2)\hat{x} + (ft^4 - et^5)\hat{y},$$

Where the constants $c, e, f,$ and h have SI units of $[c] = \left[\frac{\text{m}}{\text{s}^2}\right], [f] = \left[\frac{\text{m}}{\text{s}^4}\right], [e] = \left[\frac{\text{m}}{\text{s}^5}\right]$.

(c) Find the velocity **vector** at a later time.

(d) Find the acceleration **vector** at some later time.

[2] A ball is thrown upward at an angle of 55° with respect to the horizontal direction with an initial velocity of 5 m/s. Answer the following questions, **providing correct SI units**.

- (a)** What is the maximum height to which the ball rises?
- (b)** How long is the ball in the air?
- (c)** What is the range of the ball?
- (d)** What is the impact velocity **vector** of the ball?

[3] Consider the following vectors:

$$\vec{A}=9\hat{x}+2\hat{y}; \vec{B}=-5\hat{x}+3\hat{y}; \vec{C}=5\hat{x}-3\hat{y}$$

(a) $\vec{A} + \vec{B} + \vec{C} =$ _____


(b) $\vec{A} \cdot \vec{B} =$ _____

(c) $|\vec{A} + \vec{B}| =$ _____

(d) $(\vec{A} + \vec{C}) \cdot \hat{y} =$ _____

(e) The angle that vector A makes with respect to the positive x-axis. $\theta =$ _____

[4] Two masses resting on a frictionless table are connected by a string and forces are applied as shown. You may assume $F_1 > F_2$ here.



(a) In the boxes below, draw complete and correct free body diagrams for this system.



(b) Provide the 4 equations that come from Newton's Laws for this system.

(c) Find the acceleration of the system in terms of M_1 , M_2 , and F .

(d) Find the tension in the string in terms of M_1 , M_2 , and F .

(e) If $M_1=2\text{kg}$, $M_2=10\text{kg}$, $F=15\text{N}$, then provide numerical answers for a and T together with correct SI units.

$a =$ _____

$T =$ _____